

NORMING ALGEBRAS AND AUTOMATIC COMPLETE BOUNDEDNESS OF ISOMORPHISMS OF OPERATOR ALGEBRAS

DAVID R. PITTS

ABSTRACT. We combine the notion of norming algebra introduced by Pop, Sinclair and Smith with a result of Pisier to show that if $\mathcal{A}_1$ and $\mathcal{A}_2$ are operator algebras, then any bounded epimorphism of $\mathcal{A}_1$ onto $\mathcal{A}_2$ is completely bounded provided that $\mathcal{A}_2$ contains a norming C^* -subalgebra. We use this result to give some insights into Kadison's Similarity Problem: we show that every faithful bounded homomorphism of a C^* -algebra on a Hilbert space has completely bounded inverse, and show that a bounded representation of a C^* -algebra is similar to a $*$ -representation precisely when the image operator algebra λ -norms itself. We give two applications to isometric isomorphisms of certain operator algebras. The first is an extension of a result of Davidson and Power on isometric isomorphisms of CSL algebras. Secondly, we show that an isometric isomorphism between subalgebras $\mathcal{A}_i$ of C^* -diagonals $(\mathcal{C}_i, \mathcal{D}_i)$ ($i = 1, 2$) satisfying $\mathcal{D}_i \subseteq \mathcal{A}_i \subseteq \mathcal{C}_i$ extends uniquely to a $*$ -isomorphism of the C^* -algebras generated by $\mathcal{A}_1$ and $\mathcal{A}_2$; this generalizes results of Muhly-Qiu-Solel and Donsig-Pitts.

1. INTRODUCTION AND NORMING ALGEBRAS

Let $\mathcal{A}$ be a unital operator algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ be a homomorphism. If u is contractive (resp. isometric or bounded), it is not generally the case that u is completely contractive (resp. completely isometric or completely bounded). However, in some cases it is possible to conclude that if u is isometric or contractive, then u is completely isometric or completely contractive. For example, the contractive homomorphisms of a C^* -algebra are exactly the $*$ homomorphisms, and hence any contractive homomorphism of a C^* -algebra into $\mathcal{B}(\mathcal{H})$ is completely contractive. It is not known however if every bounded representation of a C^* -algebra is completely bounded; a result of Haagerup (stated as Theorem 2.1 below) shows this question is equivalent to Kadison's similarity problem.

The purpose of this note is to give a sufficient condition on an operator algebra $\mathcal{B}$ which ensures that every bounded epimorphism $u : \mathcal{A} \rightarrow \mathcal{B}$ of the operator algebra $\mathcal{A}$ onto $\mathcal{B}$ is completely bounded. We show that this condition can be used to give simple proofs of several results in the literature, and also that every bounded faithful representation of a C^* -algebra is bounded below.

Throughout, all operator algebras are norm-closed, and we typically use $\mathcal{A}$ and $\mathcal{B}$ to denote operator algebras. (There are several texts containing the background we need for operator spaces and operator algebras, see for example [1, 4, 12].) An *operator $\mathcal{A}$ - $\mathcal{B}$ -bimodule* is an operator space $\mathcal{M}$ which is a left- $\mathcal{A}$, right- $\mathcal{B}$ bimodule where the bimodule action is completely contractive in the sense that for any $A \in M_{np}(\mathcal{A})$, $M \in M_p(\mathcal{M})$ and $B \in M_{pn}(\mathcal{B})$,

$$\|AMB\|_{M_n(\mathcal{M})} \leq \|A\|_{M_{np}(\mathcal{A})} \|M\|_{M_p(\mathcal{M})} \|B\|_{M_{pn}(\mathcal{B})}.$$

(We shall generally not write subscripts on the norms in the sequel, unless necessary for clarity.) We will sometimes write ${}_A\mathcal{M}_B$ when $\mathcal{M}$ is an $\mathcal{A}$ - $\mathcal{B}$ -bimodule.

Date: September 20, 2006.

2000 Mathematics Subject Classification. 47L30, 46L07, 47L55.

Key words and phrases. completely bounded isomorphism, C^* -algebra, operator algebra.

For convenience, we will always assume that operator algebras are unital unless explicitly stated otherwise. However, many of the results in the sequel are valid for non-unital algebras.

Given such an $\mathcal{A}$ - $\mathcal{B}$ operator bimodule $\mathcal{M}$, we may define a family of norms η_n on $M_n(\mathcal{M})$ as follows: for $X \in M_n(\mathcal{M})$,

$$\eta_n(X) := \sup\{\|RXC\| : R \in M_{1n}(\mathcal{A}), C \in M_{n1}(\mathcal{B}) \text{ and } \max\{\|C\|, \|R\|\} \leq 1\}.$$

Clearly $\eta_n(X) \leq \|X\|_{\mathcal{M}_n(\mathcal{M})}$.

Definition 1.1. ([18]) Let $\lambda > 0$ be a real number. We say that ${}_{\mathcal{A}}\mathcal{M}_{\mathcal{B}}$ is λ -normed by $\mathcal{A}$ and $\mathcal{B}$ if for every $n \in \mathbb{N}$ and $X \in M_n(\mathcal{M})$,

$$\lambda \|X\| \leq \eta_n(X).$$

When $\lambda = 1$, we say that ${}_{\mathcal{A}}\mathcal{M}_{\mathcal{B}}$ is *normed* by $\mathcal{A}$ - $\mathcal{B}$. When $\mathcal{A} = \mathcal{B}$, we simply say that $\mathcal{M}$ is normed by $\mathcal{A}$.

When $\mathcal{A}$ and $\mathcal{B}$ are C^* -algebras, C. Pop [17] shows that $(\mathcal{M}, \{\eta_n\})$ is the smallest operator space structure on the Banach space $\mathcal{M}$ which is compatible with the module structure of $\mathcal{M}$.

We will make essential use of the following result.

Theorem 1.2 ([16, Lemma 7.7, p. 128]). *Let $\mathcal{A}$ be a C^* -algebra and suppose that $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a bounded homomorphism. Then for any $R \in M_{1,n}(\mathcal{A})$ and $C \in M_{n,1}(\mathcal{A})$ we have*

$$\begin{aligned} \|u_{1,n}(R)(u_{1,n}(R))^*\| &\leq \|u\|^4 \|RR^*\| \quad \text{and} \\ \|(u_{n,1}(C))^*u_{n,1}(C)\| &\leq \|u\|^4 \|C^*C\|. \end{aligned}$$

Remark 1.3. Actually, [16, Lemma 7.7, p. 128] gives the statement and proof for the column C . However to obtain the statement for R , one applies the statement for C to the opposite algebra $\mathcal{A}^{\text{op}}$ and the homomorphism ψ of $\mathcal{A}^{\text{op}}$ on the conjugate Hilbert space $\overline{\mathcal{H}}$ given by $d \in \mathcal{A} \mapsto \psi(d)$, where $\psi(d)\overline{h} = \overline{(u(d))^*h}$.

The following is our central observation. The key idea is to combine the techniques of [18, Theorem 2.10] with Theorem 1.2. (See also [20, Theorem 2.1].)

Theorem 1.4. *Let $\mathcal{B}$ be a norm-closed operator algebra which contains a C^* -algebra $\mathcal{D}$, and assume $\mathcal{D}$ norms $\mathcal{B}$. If $\mathcal{A}$ is a norm-closed operator algebra and $u : \mathcal{A} \rightarrow \mathcal{B}$ is a bounded isomorphism, then u is completely bounded and*

$$\|u\|_{cb} \leq \|u\| \|u^{-1}\|^4.$$

Remarks. If $\mathcal{D}$ only λ -norms $\mathcal{B}$, then u is completely bounded with $\|u\|_{cb} \leq \lambda^{-1} \|u\| \|u^{-1}\|^4$. Also, clearly the theorem applies when u is a bounded epimorphism: simply replace $\mathcal{A}$ with $\mathcal{A}/\ker u$ and u with the induced isomorphism of the quotient onto $\mathcal{B}$.

Proof. Let $T = (T_{ij}) \in M_n(\mathcal{A})$. Then for any $R \in M_{1n}(\mathcal{D})$ and $C \in M_{n1}(\mathcal{D})$ with $\|R\| \leq 1$ and $\|C\| \leq 1$ we have,

$$\begin{aligned} \|Ru_n(T)C\| &= \left\| \sum_{i,j=1}^n R_i u(T_{ij}) C_j \right\| \\ &\leq \|u\| \left\| \sum_{i,j=1}^n u^{-1}(R_i) T_{ij} u^{-1}(C_j) \right\| \\ &= \|u\| \left\| u_{1,n}^{-1}(R) T u_{n,1}^{-1}(C) \right\| \\ &\leq \|u\| \left\| u_{1,n}^{-1}(R) \right\| \|T\| \left\| u_{n,1}^{-1}(C) \right\|. \end{aligned}$$

We may assume that $\mathcal{A}$ is represented completely isometrically as operators acting on a Hilbert space $\mathcal{H}$. Thus, $u^{-1}|_{\mathcal{D}}$ is a bounded homomorphism of $\mathcal{D}$ into $\mathcal{B}(\mathcal{H})$. By Lemma 1.2, $\left\| u_{n,1}^{-1}(C) \right\| \leq \|u^{-1}\|^2$ and $\left\| u_{1,n}^{-1}(R) \right\| \leq \|u^{-1}\|^2$. Taking suprema over R and C gives

$$\|u_n(T)\| \leq \|u\| \|u^{-1}\|^4 \|T\|,$$

as desired. $\square$

When the isomorphism is isometric, more can be said. For any operator algebra $\mathcal{B}$, let $C_{\text{env}}^*(B)$ be the C^* -envelope of $\mathcal{B}$.

Corollary 1.5. *For $i = 1, 2$, suppose that $\mathcal{A}_i$ are operator algebras and $\mathcal{D}_i \subseteq \mathcal{C}_i$ is a norming C^* -subalgebra of $\mathcal{A}_i$. If $u : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ is an isometric isomorphism, then u extends uniquely to a $*$ -isomorphism $\tilde{u} : C_{\text{env}}^*(\mathcal{A}_1) \rightarrow C_{\text{env}}^*(\mathcal{A}_2)$.*

Proof. Theorem 1.4 shows that u and u^{-1} are complete contractions, so that u is a complete isometry. The result follows from the universal property of C^* -envelopes. $\square$

2. APPLICATIONS

In this section we record some consequences of Theorem 1.4. We shall require the following closely related results of Haagerup and Paulsen.

Theorem 2.1 (Haagerup [6]). *Suppose $\mathcal{A}$ is a C^* -algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a completely bounded homomorphism. Then there exists an invertible operator $S \in \mathcal{B}(\mathcal{H})$ with $\|S\| \|S^{-1}\| = \|u\|_{\text{cb}}$ such that for every $a \in \mathcal{A}$,*

$$a \mapsto Su(a)S^{-1}$$

is completely contractive (and hence a $*$ -representation).

Theorem 2.2 (Paulsen [13]). *Suppose $\mathcal{A}$ is a unital operator algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a completely bounded unital homomorphism. Then there exists an invertible operator $S \in \mathcal{B}(\mathcal{H})$ with $\|S\| \|S^{-1}\| = \|u\|_{\text{cb}}$ such that for every $a \in \mathcal{A}$,*

$$a \mapsto Su(a)S^{-1}$$

is a completely contractive homomorphism.

2.1. Applications to C^* -algebras and Kadison's Similarity Problem. We begin with a new proof of a result of Gardner.

Theorem 2.3 (Gardner [5]). *Suppose $\mathcal{A}$ and $\mathcal{B}$ are C^* -algebras and $u : \mathcal{A} \rightarrow \mathcal{B}$ is an isomorphism. Then u is (completely) bounded and there exists a $*$ -isomorphism $\alpha : \mathcal{A} \rightarrow \mathcal{B}$ and a bounded automorphism β of $\mathcal{B}$ such that $u = \beta \circ \alpha$. If $\mathcal{B} \subseteq \mathcal{B}(\mathcal{H})$, there exists a positive invertible operator $S \in \mathcal{B}(\mathcal{H})$ with $\|S\| \|S^{-1}\| \leq \|u\|$ so that $\beta = \text{Ad } S$.*

Proof. By [18, Lemma 2.3(i)], $\mathcal{B}$ norms itself, and as observed by Gardner, u is bounded. (One can also use a result of B. Johnson [7] (see also [19]) concerning automatic continuity of a homomorphism from a Banach algebra onto a semi-simple Banach algebra.) Theorem 1.4 implies that u is completely bounded. Assume that $\mathcal{B} \subseteq \mathcal{B}(\mathcal{H})$. By Theorem 2.1, there exists an invertible operator T with $\|T\| \|T^{-1}\| = \|u\|_{\text{cb}}$ such that $\text{Ad } T \circ u$ is a completely contractive homomorphism, so that $\text{Ad } T \circ u$ is therefore a $*$ -homomorphism. Let $S = |T|$, and let U be the polar part of T , so $T = US$. Then $(\text{Ad } S)(\mathcal{B})$ is a C^* -algebra, and it follows that $S^2 \mathcal{B} S^{-2} = \mathcal{B}$. By Gardner's Invariance Theorem [5, Theorem 3.5], $\beta := \text{Ad } S^{-1}$ is an automorphism of $\mathcal{B}$. Now let $\alpha = \text{Ad}(U^*T) \circ u$. Since $\text{Ad } T \circ u$ is a $*$ -isomorphism of $\mathcal{A}$ onto its range, α is a $*$ -isomorphism of $\mathcal{A}$ onto $\mathcal{B}$. Then $u = \beta \circ \alpha$. $\square$

Remark 2.4. Gardner's original arguments give somewhat more. In particular, he shows that if $\mathcal{A}$ and $\mathcal{B}$ are faithfully represented using the universal atomic representations, then α can be taken to have the form $\text{Ad } U$ for some unitary U .

The following is an immediate corollary of Theorem 1.4.

Theorem 2.5. *Suppose $\mathcal{A}$ is an operator algebra and $\mathcal{B}$ is a C^* -algebra. If $u : \mathcal{A} \rightarrow \mathcal{B}$ is an isomorphism, then u is automatically completely bounded and $\|u\|_{\text{cb}} \leq \|u\| \|u^{-1}\|^4$.*

Proof. Continuity of u follows from Johnson's theorem [7], then complete boundedness follows from Theorem 1.4 together with the fact that a C^* -algebra norms itself [18, Lemma 2.3(i)]. $\square$

We now wish to make some observations regarding Kadison's Similarity Problem. Recall that this problem asks whether every bounded representation of a C^* -algebra is similar to a $*$ -representation, which as noted above, is equivalent to the question of whether bounded representations of C^* -algebras are automatically completely bounded. Theorem 2.5 can be used to prove that bounded representations of C^* -algebras are (modulo the kernel) "completely bounded below."

Theorem 2.6. *Suppose $\mathcal{A}$ is a C^* -algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a bounded homomorphism. Put $\mathcal{J} = \ker u$. Then there exists a real number $k > 0$ such that for every $n \in \mathbb{N}$ and every $T \in M_n(\mathcal{A})$,*

$$k \text{dist}(T, M_n(\mathcal{J})) \leq \|u_n(T)\|.$$

Proof. Since $\mathcal{A}/\mathcal{J}$ is a C^* -algebra isomorphic under the map induced by u to $u(\mathcal{A})$, without loss of generality we may assume that u is one-to-one. Thus our task is to prove that $u := u^{-1}$ is completely bounded. This will follow from Theorem 2.5 once we prove that the image $\mathcal{B} := u(\mathcal{A})$ is closed, and hence an operator algebra. We may also assume that $\mathcal{A}$ is unital and that $u(I) = I$.

Let x be a non-zero element of $\mathcal{A}$, and let $\mathcal{C}$ be the unital C^* -subalgebra of $\mathcal{A}$ generated by x^*x . Since $\mathcal{C}$ is abelian, the restriction of u to $\mathcal{C}$ is completely bounded on $\mathcal{C}$, so by the Dixmier-Day Theorem on amenable groups, there exists an invertible operator S with $\|S\| \|S^{-1}\| \leq \|u|_{\mathcal{C}}\|^2$ such that $(\text{Ad } S) \circ u|_{\mathcal{C}}$ is a $*$ -homomorphism. Since u is one-to-one, we have

$$\|x\|^2 = \|x^*x\| = \|Su(x^*x)S^{-1}\| \leq \|u\|^2 \|u(x^*)u(x)\| \leq \|u\|^3 \|x\| \|u(x)\|,$$

so that $\|x\| \leq \|u\|^3 \|u(x)\|$. Thus, u is bounded below, so that the range of u is closed. $\square$

Combining Theorems 2.2 and 2.6 with the structure of completely contractive representations we obtain the following corollary.

Corollary 2.7. *Suppose $\mathcal{A} \subseteq \mathcal{B}(\mathcal{H})$ is a C^* -algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H}_u)$ is a faithful bounded representation. Then there exists a Hilbert space $\mathcal{K}$, a $*$ -representation $\pi : \mathcal{B}(\mathcal{H}_u) \rightarrow \mathcal{B}(\mathcal{K})$, an isometry $W : \mathcal{H} \rightarrow \mathcal{K}$ and an invertible operator $S \in \mathcal{B}(\mathcal{H})$ with $\|S\| \|S^{-1}\| \leq \|u^{-1}\| \|u\|^4$ such that for every $x \in \mathcal{A}$,*

$$(1) \quad x = SW^* \pi(u(x)) W S^{-1}.$$

Proof. Let $\mathcal{B} = u(\mathcal{A})$. We may assume that $\mathcal{A}$ is unital and $u(I) = I$. Theorem 2.6 shows that u^{-1} is a completely bounded map from $\mathcal{B}$ to $\mathcal{B}(\mathcal{H})$. Therefore, by Theorem 2.2, there exists an invertible operator $S \in \mathcal{B}(\mathcal{H})$ with $\|S\| \|S^{-1}\| = \|u^{-1}\|_{cb}$ so that $\psi := \text{Ad}(S^{-1}) \circ u^{-1}$ is completely contractive. By Arveson's Structure Theorem for completely contractive representations of operator algebras, there exists a Hilbert space $\mathcal{K}$, a $*$ -representation $\pi : \mathcal{B}(\mathcal{H}_u) \rightarrow \mathcal{B}(\mathcal{K})$ and an isometry W so that for all $b \in \mathcal{B}$, $\psi(b) = W^* \pi(b) W$. Letting $b = u(x)$ (for $x \in \mathcal{A}$) yields (1). The estimate for the condition number of S follows from Theorem 1.4. $\square$

Remark 2.8. Unfortunately, we have been unable to solve for $u(x)$ in (1); doing so would of course lead to a solution of Kadison's problem.

The range of the isometry W appearing in Corollary 1 is a semi-invariant subspace for $\pi(\mathcal{B})$. Thus, $WW^* = PQ^\perp$ for some projections $P, Q \in \text{Lat}(\pi(\mathcal{B}))$, with $Q \leq P$. The map $x \mapsto WS^{-1}xSW^*$ is a homomorphism into the “2,2-diagonal piece” of $\pi(u(x))$ relative to the block decomposition of $\pi(u(x))$ according to $I = Q + PQ^\perp + Q^\perp P^\perp$.

We now show that Kadison problem is equivalent to the issue of whether the image of a C^* -algebra under a bounded homomorphism λ norms itself.

Theorem 2.9. *Suppose $\mathcal{A}$ is a C^* -algebra and $u : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is a bounded representation with image $\mathcal{B} := u(\mathcal{A})$, and let $\tilde{u} : \mathcal{A}/\ker u \rightarrow \mathcal{B}$ be the induced map. If $\mathcal{B}$ λ -norms itself for some $\lambda > 0$, then u is completely bounded and $\|u\|_{cb} \leq \lambda^{-1} \|u\|^9 \|\tilde{u}^{-1}\|^2$. Conversely, if u is completely bounded, then $\mathcal{B}$ λ -norms itself for any λ with $0 < \lambda \leq \frac{1}{\|u\|_{cb} \|\tilde{u}^{-1}\| \|u\|^4}$.*

Proof. We may assume that u is a monomorphism, so Theorem 2.5 gives $\|u^{-1}\|_{cb} \leq \|u\|^4 \|u^{-1}\|$. Suppose that $\mathcal{B}$ is λ -normed by itself for some $\lambda > 0$, and let $T \in M_n(\mathcal{A})$. The calculation in the proof of Theorem 1.4 shows that if $R \in M_{1,n}(\mathcal{B})$ and $C \in M_{n,1}(\mathcal{B})$ satisfy $\|R\| \leq 1$ and $\|C\| \leq 1$, then

$$\|Ru_n(T)C\| \leq \|u\| \left\| u_{1,n}^{-1}(R) \right\| \|T\| \left\| u_{n,1}^{-1}(C) \right\| \leq \|u\|^9 \|u^{-1}\|^2 \|T\|.$$

Taking suprema over all such R and C gives

$$\lambda \|u_n(T)\| \leq \|u\|^9 \|u^{-1}\|^2 \|T\|,$$

so u is completely bounded with $\|u\|_{cb} \leq \lambda^{-1} \|u\|^9 \|u^{-1}\|^2$.

Conversely, suppose that u is completely bounded, and view $\mathcal{B}$ as a bimodule over itself. For any $R \in \mathcal{M}_{1,n}(\mathcal{A})$, $C \in \mathcal{M}_{n,1}(\mathcal{A})$ with $\|R\|, \|C\| \leq 1$, we have,

$$\begin{aligned} \|Ru_n^{-1}(T)C\| &\leq \|u^{-1}\| \|u(Ru_n^{-1}(T)C)\| \\ &= \|u^{-1}\| \|u_{1,n}(R)\| \|u_{n,1}(C)\| \left\| \frac{u_{1,n}(R)}{\|u_{1,n}(R)\|} T \frac{u_{n,1}(C)}{\|u_{n,1}(C)\|} \right\| \\ &\leq \|u^{-1}\| \|u\|^4 \eta_n(T) \quad (\text{using Lemma 1.2}). \end{aligned}$$

Taking suprema yields $\|u_n^{-1}(T)\| \leq \|u^{-1}\| \|u\|^4 \eta_n(T)$, and hence

$$\|T\| \leq \|u_n\| \|u_n^{-1}(T)\| \leq \|u\|_{cb} \|u^{-1}\| \|u\|^4 \eta_n(T).$$

Thus $\mathcal{B}$ λ -norms itself for any $\lambda \leq \frac{1}{\|u\|_{cb} \|u^{-1}\| \|u\|^4}$. $\square$

Corollary 2.10. *Suppose $\mathcal{B} \subseteq \mathcal{B}(\mathcal{H})$ is an operator algebra which is isomorphic to a C^* -algebra A . Then there exists an invertible operator $S \in \mathcal{B}(\mathcal{H})$ such that $S\mathcal{B}S^{-1}$ is a C^* -algebra if and only if $\mathcal{B}$ λ -norms itself for some $\lambda > 0$.*

Proof. Let $u : \mathcal{A} \rightarrow \mathcal{B}$ be the isomorphism. Then Theorem 2.9 shows that if $\mathcal{B}$ λ -norms itself, then u is completely bounded, hence by Theorem 2.1, u is similar to a $*$ -representation, and so $\mathcal{B}$ is similar to a C^* -algebra. Conversely, if $\mathcal{B}$ is similar to a C^* -algebra, then u is similar to a $*$ -representation and an application of Theorem 2.9 completes the proof. $\square$

The following question is thus a reformulation of Kadison's question.

Question 2.11. Suppose $\mathcal{B}$ is an operator algebra which is isomorphic to a C^* -algebra. Does $\mathcal{B}$ λ -norm itself for some $\lambda > 0$?

Remark 2.12. Haagerup [6] showed that every bounded, cyclic representation u of a C^* -algebra $\mathcal{A}$ is completely bounded with $\|u\|_{cb} \leq \|u\|^3$. So given an arbitrary representation u of $\mathcal{A}$ on $\mathcal{H}$, let $\mathcal{B} = u(\mathcal{A})$. For each unit vector $\xi \in \mathcal{H}$, let P_ξ be the projection onto the cyclic subspace $[\mathcal{B}\xi]$ and let $\mathcal{B}_\xi$ be the restriction of $\mathcal{B}$ to $\mathcal{H}_\xi := P_\xi \mathcal{H}$. The representation u_ξ of $\mathcal{A}$ given by $u_\xi(x) = u(x)P_\xi$ is thus completely bounded and $\|u_\xi\|_{cb} \leq \|u\|_{cb}$. Theorem 2.2 shows that there exists an invertible operator $S_\xi \in \mathcal{B}(\mathcal{H}_\xi)$ such that $\|u_\xi\|_{cb} = \|S_\xi\| \|S_\xi^{-1}\|$ and $(\text{Ad } S_\xi)(\mathcal{B}_\xi)$ is a C^* -algebra, call it $\mathcal{A}_\xi$. Applying Theorem 2.9 to $\text{Ad } S_\xi^{-1} : \mathcal{A}_\xi \rightarrow \mathcal{B}_\xi$, we find $\|\text{Ad } S_\xi\| \leq \|S_\xi\| \|S_\xi^{-1}\|$, so that $\mathcal{B}_\xi$ λ -norms itself with $\lambda = \|u\|^{-18}$.

In the following example, we show that there exists an operator algebra which does not λ -norm itself. The idea for the proof is due to Ken Davidson.

Example 2.13. Let $\mathbb{D} \subseteq \mathbb{C}$ be the open unit disk and let $\mathbb{A}(\mathbb{D}) \subseteq C(\overline{\mathbb{D}})$ be the disk algebra, that is, the collection of all continuous functions on the closed unit disk which are analytic in $\mathbb{D}$. We use $\mathcal{P} \subseteq \mathbb{A}(\mathbb{D})$ to denote the collection of all polynomials.

Recall that an operator $T \in \mathcal{B}(\mathcal{H})$ is polynomially bounded if there exists $K > 0$ such that for every $p \in \mathcal{P}$, $\|p(T)\| \leq K \|p\|$. Polynomial boundedness of T is equivalent to the existence of a bounded homomorphism $u : \mathbb{A}(\mathbb{D}) \rightarrow \mathcal{B}(\mathcal{H})$ such that for every $p \in \mathcal{P}$, $u(p) = p(T)$. A polynomially bounded operator T is *completely polynomially bounded* if u is completely bounded. Paulsen [14] showed that T is completely polynomially bounded if and only if T is similar to a contraction. Pisier [15] showed that there exists a polynomially bounded operator $T \in \mathcal{B}(\mathcal{H})$ which is not completely polynomially bounded, so T is not similar to a contraction.

Fix a polynomially bounded operator $T \in \mathcal{B}(\mathcal{H})$. Notice that the spectrum of T is contained in $\overline{\mathbb{D}}$. If U is a unitary operator with $\sigma(U) = \mathbb{T}$, then T is completely polynomially bounded if and only if $T \oplus U$ is completely polynomially bounded. We may therefore assume that

$$\mathbb{T} \subseteq \sigma(T).$$

Let $u : \mathbb{A}(\mathbb{D}) \rightarrow \mathcal{B}(\mathcal{H})$ be the extension of the map $p \in \mathcal{P} \mapsto p(T)$ and set $\mathcal{B} = u(\mathbb{A}(\mathbb{D}))$. Since $\mathbb{T} \subseteq \sigma(T)$, we have $\|u(f)\| \geq \|f\|$ for every $f \in \mathbb{A}(\mathbb{D})$, so that $\mathcal{B}$ is closed and u^{-1} is contractive. Since $\mathbb{A}(\mathbb{D}) \subseteq C(\overline{\mathbb{D}})$, the operator space structure on $\mathbb{A}(\mathbb{D})$ is minimal among all all possible operator space structures on $\mathbb{A}(\mathbb{D})$ (see [1, Paragraph 1.2.21]), hence u^{-1} is completely contractive.

View $\mathcal{B}$ as a $\mathcal{B}$ -bimodule and let η_n be the norm on $M_n(\mathcal{B})$ as in Definition 1.1. We shall show that for every n , the norm $\eta_n(u_n(\cdot))$ and the usual norm $\|\cdot\|_{M_n(\mathbb{A}(\mathbb{D}))}$ are equivalent norms on $M_n(\mathbb{A}(\mathbb{D}))$.

Choose $X \in M_n(\mathbb{A}(\mathbb{D}))$, $R \in M_{1n}(\mathbb{A}(\mathbb{D}))$ and $C \in M_{n1}(\mathbb{A}(\mathbb{D}))$. Since u is bounded, we have $\|u_{1n}(R)u_n(X)u_{n1}(C)\| = \|u(RXC)\| \leq \|u\| \|RXC\|$. Since $\|u_{1n}(R)\| \geq \|R\|$ and $\|u_{n1}(C)\| \geq \|C\|$, we obtain

$$(2) \quad \eta_n(u_n(X)) \leq \|u\| \|X\|.$$

On the other hand, a result of Bourgain [2] (see also [16, Theorem 9.9]) shows that there exists a constant $s > 0$ (independent of u) such that for every n , $\max\{\|u_{1n}\|, \|u_{n1}\|\} \leq s \|u\|$. Therefore,

$$\begin{aligned} \|RXC\| &\leq \|u_{1n}(R)u_n(X)u_{n1}(C)\| \\ &\leq s^2 \|u\|^2 \left\| \frac{u_{1n}(R)}{\|u_{1n}(R)\|} u_n(X) \frac{u_{n1}(C)}{\|u_{n1}(C)\|} \right\| \|R\| \|C\|. \end{aligned}$$

Since the operator space structure on $\mathbb{A}(\mathbb{D})$ is minimal, $\mathbb{A}(\mathbb{D})$ norms itself. Taking the supremum over R and C with norm one, we obtain

$$(3) \quad \|X\| \leq s^2 \|u\|^2 \eta_n(u_n(X)).$$

Combining (2) and (3) establishes the claim.

Thus, when T is chosen to be polynomially bounded but not completely polynomially bounded, we see that $\mathcal{B}$ cannot λ -norm itself.

2.2. An Application to CSL Algebras. Recall that a *CSL algebra* is an operator algebra $\mathcal{A} \subseteq \mathcal{B}(\mathcal{H})$ which is both reflexive and such that there exists a MASA $\mathcal{D} \subseteq \mathcal{B}(\mathcal{H})$ with $\mathcal{D} \subseteq \mathcal{A}$. The lattice of invariant projections of a CSL algebra is a commutative family of projections, and when this lattice is completely distributive, the CSL algebra is called a *completely distributive* CSL algebra.

Davidson and Power proved that isometric isomorphisms of completely distributive CSL algebras are unitarily implemented. Their techniques involved homological ideas and were somewhat intricate. We can give a simpler proof of their result, and which also extends theirs. For convenience, we assume irreducibility. If $\mathcal{A}$ is not irreducible, one can use direct integrals along the center of $\mathcal{A} \cap \mathcal{A}^*$, together with an appropriate hypothesis on the lattices of each term in the direct integral to obtain a more general result.

Theorem 2.14. *For $i = 1, 2$, let $\mathcal{A}_i \subseteq \mathcal{B}(\mathcal{H}_i)$ be CSL algebras such that $\mathcal{A}_i \cap \mathcal{K}_i \neq (0)$ and suppose that $\mathcal{A}_1$ is irreducible. If $u : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ is an isometric isomorphism, then there exists a unitary operator $U \in \mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ such that $u = \text{Ad } U$.*

Proof. Since the isomorphism u is isometric, its restriction to $\mathcal{A}_1 \cap (\mathcal{A}_1)^*$ is a $*$ -isomorphism, so in particular, $\text{Lat}(\mathcal{A}_2)$ is isomorphic to $\text{Lat}(\mathcal{A}_1)$. Thus, since $\mathcal{A}_1$ is irreducible, so is $\mathcal{A}_2$. Let $\mathcal{C}_i$ be the C^* -subalgebra of $\mathcal{B}(\mathcal{H}_i)$ generated by $\mathcal{A}_i$. Then $\mathcal{C}_i$ is an irreducible C^* -algebra containing a compact operator, hence $\mathcal{C}_i$ contains all the compact operators. There exists a $*$ -epimorphism $\pi : \mathcal{C}_i \rightarrow C_{\text{env}}^*(\mathcal{A}_i)$ such that $\pi|_{\mathcal{A}_i} = \iota$, where ι is the canonical inclusion of $\mathcal{A}_i$ into $C_{\text{env}}^*(\mathcal{A}_i)$. If $\ker \pi \neq (0)$, then since $\mathcal{C}_i$ contains the compact operators, $\ker \pi \cap \mathcal{K}_i \neq (0)$. Hence $\mathcal{K}_i \subseteq \ker \pi$, since $\ker \pi \cap \mathcal{K}_i$ is an ideal in an irreducible C^* -algebra. But this is impossible, since π is isometric on $\mathcal{A}_i \cap \mathcal{K}_i$. Therefore $\ker \pi = (0)$, so that $\mathcal{C}_i$ is the C^* -envelope of $\mathcal{A}_i$.

Theorem 2.7 of [18] shows that any MASA is norming for $\mathcal{B}(\mathcal{H})$, hence $\mathcal{A}_i$ contain norming C^* -subalgebras. Theorem 1.4 shows that u and u^{-1} are completely contractive, so that u is a complete isometry. By the universal property of C^* -envelopes (applied to u and u^{-1}), u extends to a $*$ -isomorphism $\tilde{u}$ of $\mathcal{C}_1$ onto $\mathcal{C}_2$. The compact operators are the smallest closed two-sided ideal contained in $\mathcal{C}_i$, so that the restriction of $\tilde{u}$ to the compact operators is a $*$ -isomorphism of $\mathcal{K}(\mathcal{H}_1)$ onto $\mathcal{K}(\mathcal{H}_2)$. Therefore, there exists a unitary operator U so that $(\text{Ad } U)|_{\mathcal{K}(\mathcal{H}_1)} = \tilde{u}|_{\mathcal{K}(\mathcal{H}_1)}$. Finally,

if $T \in \mathcal{C}_1$ and if $\eta \in \mathcal{H}_2$, we may find a finite rank projection P so that $2\|T\|\|P^\perp\eta\| < \varepsilon$. Then since $\tilde{u}^{-1}(P) = (\text{Ad } U^*)(P)$, we have

$$\begin{aligned} \|(\tilde{u}(T) - (\text{Ad } U)(T))\eta\| &\leq \|(\tilde{u}(T\tilde{u}^{-1}(P)) - (\text{Ad } U)(T(\text{Ad } U^*(P))))\eta\| \\ &\quad + \|(\tilde{u}(T) - (\text{Ad } U)(T))P^\perp\eta\| \\ &= \|(\tilde{u}(T) - (\text{Ad } U)(T))P^\perp\eta\| < \varepsilon, \end{aligned}$$

so $\tilde{u}(T) = (\text{Ad } U)(T)$. Since $\mathcal{A}_1 \subseteq \mathcal{C}_1$, the proof is complete. $\square$

2.3. Applications to Subalgebras of C^* -Diagonals. In this subsection, we provide applications to subalgebras of certain classes of C^* -algebras.

A C^* -diagonal is a pair $(\mathcal{C}, \mathcal{D})$ of C^* -algebras such that $\mathcal{D}$ is abelian and such that

- i) every pure state of $\mathcal{D}$ extends uniquely to a pure state of $\mathcal{C}$;
- ii) the conditional expectation $E : \mathcal{C} \rightarrow \mathcal{D}$ (whose existence is guaranteed by (i)) is faithful;
- iii) the closed linear span of the set $\{v \in \mathcal{C} : v\mathcal{D} = \mathcal{D}v\}$ is $\mathcal{C}$.

We will assume that both $\mathcal{C}$ and $\mathcal{D}$ are unital. The extension property then implies that $\mathcal{D}$ is a MASA in $\mathcal{C}$.

Such pairs were introduced by Kumjian [8], who used slightly different, but essentially equivalent axioms (see [3] for a discussion of the equivalence). Also, C^* -diagonals and their subalgebras were further in several papers, see for example [3, 10, 11].

Our first task is to show that $\mathcal{D}$ norms $\mathcal{C}$.

Lemma 2.15. *Suppose $(\mathcal{C}, \mathcal{D})$ is a C^* -diagonal. Then $\mathcal{C}$ is normed by $\mathcal{D}$.*

Proof. Theorem 5.9 of [3] shows that there exists a faithful $*$ -representation $\pi : \mathcal{C} \rightarrow \mathcal{B}(\mathcal{H})$ such that $\pi(\mathcal{D})''$ is a MASA in $\mathcal{B}(\mathcal{H})$. It follows from [18, Lemma 2.2 and Theorem 2.7] that $\pi(\mathcal{D})$ norms $\mathcal{B}(\mathcal{H})$, hence $\pi(\mathcal{D})$ norms $\pi(\mathcal{C})$. As π is a faithful $*$ -representation of a C^* -algebra, it is a complete isometry, so $\mathcal{D}$ norms $\mathcal{C}$. $\square$

The following notation will be useful. When $(\mathcal{C}, \mathcal{D})$ is a C^* -diagonal and $\mathcal{A}$ is a norm closed algebra with $\mathcal{D} \subseteq \mathcal{A} \subseteq \mathcal{C}$, we will write $\mathcal{A} \subseteq (\mathcal{C}, \mathcal{D})$.

For $i = 1, 2$, let $(\mathcal{C}_i, \mathcal{D}_i)$ be C^* -diagonals. Muhly, Qiu and Solel [11, Theorem 1.1] proved that when $\mathcal{A}_i \subseteq (\mathcal{C}_i, \mathcal{D}_i)$ are triangular, that is $\mathcal{A}_i \cap (\mathcal{A}_i)^* = \mathcal{D}_i$, which generate $\mathcal{C}_i$ and $(\mathcal{C}_i, \mathcal{D}_i)$ are nuclear, then an isometric isomorphism $u : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ extends to a $*$ -isomorphism of $\mathcal{C}_1$ onto $\mathcal{C}_2$. Later Donsig and Pitts [3, Theorem 8.9] extended this result: they showed that the hypothesis of nuclearity can be removed. To prove their result, Donsig and Pitts used showed that the isometric isomorphism between $\mathcal{A}_1$ and $\mathcal{A}_2$ induces isomorphism of an appropriate CSL algebras, then used the structure theory for isomorphisms of CSL algebras. The techniques used to prove [11, Theorem 1.1] and [3, Theorem 8.9] do not apply for non-triangular subalgebras. A

Donsig and Pitts [3, Theorem 4.22] showed that the C^* -envelope of any subalgebra (triangular or not) $\mathcal{A} \subseteq (\mathcal{C}, \mathcal{D})$ is the C^* -subalgebra, $C^*(\mathcal{A})$, of $\mathcal{C}$ generated by $\mathcal{A}$. In the context of both [11, Theorem 1.1] and [3, Theorem 8.9], $(C^*(\mathcal{A}_i), \mathcal{D}_i)$ are C^* -diagonals. When $\mathcal{C}$ is separable and nuclear, the Spectral Theorem for Bimodules [10] shows that $(C^*(\mathcal{A}), \mathcal{D})$ is a C^* -diagonal. In the general case however, it is not clear that the pair $(C^*(\mathcal{A}), \mathcal{D})$ is a C^* -diagonal—one needs to verify that condition (ii) of the definition of C^* -diagonal holds. The following consequence of [3, Theorem 4.22], Corollary 1.5 and Lemma 2.15 is therefore a significant extension of [11, Theorem 1.1] and [3, Theorem 8.9].

Theorem 2.16. *Let $\mathcal{A}_i \subseteq (\mathcal{C}_i, \mathcal{D}_i)$ be norm-closed subalgebras of C^* -diagonals. If $u : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ is an isometric isomorphism, then u extends uniquely to a $*$ -isomorphism of $C^*(\mathcal{A}_1)$ onto $C^*(\mathcal{A}_2)$.*

Remark 2.17. In [9], Mercer proves a result similar to Theorem 2.16, but where the algebras $\mathcal{A}_i$ are taken to be weak-* closed subalgebras of von Neumann algebras $\mathfrak{M}_i$ and there are Cartan MASAs $\mathcal{D}_i \subseteq \mathfrak{M}_i$ such that $\mathcal{D}_i \subseteq \mathcal{A}_i \subset \mathfrak{M}_i$. We expect that Cartan MASAs norm their containing von Neumann algebras, and thus expect that it should be possible to give a proof of Mercer’s result based on Theorem 1.4 as well.

ACKNOWLEDGMENT: This work was completed during a visit to the University of Waterloo. The author thanks Ken Davidson and the University of Waterloo for their hospitality, and also thanks Ken Davidson and Allan Donsig for several helpful conversations and suggestions.

REFERENCES

1. David P. Blecher and Christian Le Merdy, *Operator algebras and their modules—an operator space approach*, London Mathematical Society Monographs. New Series, vol. 30, The Clarendon Press Oxford University Press, Oxford, 2004, Oxford Science Publications. MR2111973
2. J. Bourgain, *New Banach space properties of the disc algebra and H^∞* , Acta Math. **152** (1984), no. 1-2, 1–48. MR736210 (85j:46091)
3. Allan P. Donsig and David R. Pitts, *Coordinate systems and bounded isomorphisms*, J. Operator Theory, to appear; arXiv:math.OA/0506627.
4. Edward G. Effros and Zhong-Jin Ruan, *Operator spaces*, London Mathematical Society Monographs. New Series, vol. 23, The Clarendon Press Oxford University Press, New York, 2000. MR1793753 (2002a:46082)
5. L. Terrell Gardner, *On isomorphisms of C^* -algebras*, Amer. J. Math. **87** (1965), 384–396. MR0179637 (31 #3883)
6. Uffe Haagerup, *Solution of the similarity problem for cyclic representations of C^* -algebras*, Ann. of Math. (2) **118** (1983), no. 2, 215–240. MR717823 (85d:46080)
7. B. E. Johnson, *The uniqueness of the (complete) norm topology*, Bull. Amer. Math. Soc. **73** (1967), 537–539. MR0211260 (35 #2142)
8. Alexander Kumjian, *On C^* -diagonals*, Canad. J. Math. **38** (1986), no. 4, 969–1008. 88a:46060
9. Richard Mercer, *Isometric isomorphisms of Cartan bimodule algebras*, J. Funct. Anal. **101** (1991), no. 1, 10–24. MR1132304 (92k:46102)
10. Paul S. Muhly, Chao Xin Qiu, and Baruch Solel, *Coordinates, nuclearity and spectral subspaces in operator algebras*, J. Operator Theory **26** (1991), no. 2, 313–332. 94i:46075
11. ———, *On isometries of operator algebras*, J. Funct. Anal. **119** (1994), no. 1, 138–170. 95a:46080
12. Vern Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics, vol. 78, Cambridge University Press, Cambridge, 2002. MR1976867 (2004c:46118)
13. Vern I. Paulsen, *Completely bounded homomorphisms of operator algebras*, Proc. Amer. Math. Soc. **92** (1984), no. 2, 225–228. 85m:47049
14. ———, *Every completely polynomially bounded operator is similar to a contraction*, J. Funct. Anal. **55** (1984), no. 1, 1–17. 86c:47021
15. Gilles Pisier, *A polynomially bounded operator on Hilbert space which is not similar to a contraction*, J. Amer. Math. Soc. **10** (1997), no. 2, 351–369. MR1415321 (97f:47002)
16. ———, *Similarity problems and completely bounded maps*, expanded ed., Lecture Notes in Mathematics, vol. 1618, Springer-Verlag, Berlin, 2001, Includes the solution to “The Halmos problem”. MR1818047 (2001m:47002)
17. Ciprian Pop, *Bimodules normés représentables sur des espaces hilbertiens*, Operator theoretical methods (Timișoara, 1998), Theta Found., Bucharest, 2000, pp. 331–370. MR1770332 (2001h:46111)
18. Florin Pop, Allan M. Sinclair, and Roger R. Smith, *Norming C^* -algebras by C^* -subalgebras*, J. Funct. Anal. **175** (2000), no. 1, 168–196. 2001h:46105
19. Allan M. Sinclair, *Automatic continuity of linear operators*, Cambridge University Press, Cambridge, 1976, London Mathematical Society Lecture Note Series, No. 21. MR0487371 (58 #7011)
20. R. R. Smith, *Completely bounded module maps and the Haagerup tensor product*, J. Funct. Anal. **102** (1991), no. 1, 156–175. MR1138841 (93a:46115)

DEPT. OF MATHEMATICS, UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE, 68588-0130
E-mail address: dpitts2@math.unl.edu